



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2015

Bridging to transplant with azacitidine in juvenile myelomonocytic leukemia: a retrospective analysis of the EWOG-MDS study group

Cseh, Annamaria ; Niemeyer, Charlotte M ; Yoshimi, Ayami ; Dworzak, Michael ; Hasle, Henrik ; van den Heuvel-Eibrink, Marry M ; Locatelli, Franco ; Masetti, Riccardo ; Schmugge, Markus ; Groß-Wieltsch, Ute ; Candás, Andrea ; Kulozik, Andreas E ; Olcay, Lale ; Suttorp, Meinolf ; Furlan, Ingrid ; Strahm, Brigitte ; Flotho, Christian

DOI: <https://doi.org/10.1182/blood-2015-01-619734>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-119486>

Journal Article

Accepted Version

Originally published at:

Cseh, Annamaria; Niemeyer, Charlotte M; Yoshimi, Ayami; Dworzak, Michael; Hasle, Henrik; van den Heuvel-Eibrink, Marry M; Locatelli, Franco; Masetti, Riccardo; Schmugge, Markus; Groß-Wieltsch, Ute; Candás, Andrea; Kulozik, Andreas E; Olcay, Lale; Suttorp, Meinolf; Furlan, Ingrid; Strahm, Brigitte; Flotho, Christian (2015). Bridging to transplant with azacitidine in juvenile myelomonocytic leukemia: a retrospective analysis of the EWOG-MDS study group. *Blood*, 125(14):2311-2313.

DOI: <https://doi.org/10.1182/blood-2015-01-619734>

Contribution: I.B., B.M., and J.S. performed molecular analysis and analyzed data; P.K. contributed vital patient information; T.H. and O.W. performed genetic analysis; L.Q.-M. and F.F. designed the study, performed the histologic analysis, analyzed data, and wrote the paper.

F.F. and L.Q.-M. contributed equally to this work.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

Correspondence: Leticia Quintanilla-Martinez, Institute of Pathology, University Hospital Tübingen, Liebermeisterstrasse 8, 72076 Tübingen, Germany; e-mail: leticia.quintanilla-fend@med.uni-tuebingen.de.

References

- Cazzola M, Kralovics R. From Janus kinase 2 to calreticulin: the clinically relevant genomic landscape of myeloproliferative neoplasms. *Blood*. 2014; 123(24):3714-3719.
- Klampfl T, Gisslinger H, Harutyunyan AS, et al. Somatic mutations of calreticulin in myeloproliferative neoplasms. *N Engl J Med*. 2013;369(25): 2379-2390.
- Nangalia J, Massie CE, Baxter EJ, et al. Somatic CALR mutations in myeloproliferative neoplasms with nonmutated JAK2. *N Engl J Med*. 2013; 369(25):2391-2405.
- Rumi E, Pietra D, Ferretti V, et al; Associazione Italiana per la Ricerca sul Cancro Gruppo Italiano Malattie Mieloproliferative Investigators. JAK2 or CALR mutation status defines subtypes of essential thrombocythemia with substantially different clinical course and outcomes. *Blood*. 2014;123(10): 1544-1551.
- Rumi E, Harutyunyan AS, Pietra D, et al; Associazione Italiana per la Ricerca sul Cancro Gruppo Italiano Malattie Mieloproliferative Investigators. CALR exon 9 mutations are somatically acquired events in familial cases of essential thrombocythemia or primary myelofibrosis. *Blood*. 2014;123(15): 2416-2419.
- Lundberg P, Nienhold R, Ambrosetti A, Cervantes F, Pérez-Encinas MM, Skoda RC. Somatic mutations in calreticulin can be found in pedigrees with familial predisposition to myeloproliferative neoplasms. *Blood*. 2014;123(17): 2744-2745.
- Hussein K, Bock O, Seegers A, et al. Myelofibrosis evolving during imatinib treatment of a chronic myeloproliferative disease with coexisting BCR-ABL translocation and JAK2V617F mutation. *Blood*. 2007;109(9): 4106-4107.
- Yamada O, Mahfoudhi E, Plo I, et al. Emergence of a BCR-ABL translocation in a patient with the JAK2V617F mutation: evidence for secondary acquisition of BCR-ABL in the JAK2V617F clone. *J Clin Oncol*. 2014;32(21): e76-e79.
- Bocchia M, Vannucchi AM, Gozzetti A, et al. Insights into JAK2-V617F mutation in CML. *Lancet Oncol*. 2007;8(10):864-866.

© 2015 by The American Society of Hematology

To the editor:

Bridging to transplant with azacitidine in juvenile myelomonocytic leukemia: a retrospective analysis of the EWOG-MDS study group

DNA methyltransferase-inhibiting azanucleosides have become a mainstay of treatment of myeloid neoplasms in adult patients,¹ with 5-azacytidine (azacitidine) being the agent in broadest clinical use. Although not curative, treatment with azacitidine achieves hematologic improvement and transfusion independency in many patients and prolongs survival.^{2,3} Even though most children with juvenile myelomonocytic leukemia (JMML) qualify for allogeneic hematopoietic stem cell transplantation (HSCT), the acceptable toxicity of low-dose azacitidine and its cytoreductive potential make it an attractive option as a bridging therapy before HSCT⁴ or as palliation after 1 or more transplants have failed. We previously published the first case report of a boy with JMML who achieved a complete clinical and genetic remission after 8 cycles of azacitidine.⁵ Here, we present a retrospective compilation of 12 children with JMML who received individual off-label treatment with azacitidine before HSCT (N = 9) or for relapsed disease (N = 3).

The children were treated at 11 centers November 2007-April 2012. Ten children were enrolled in the studies “98” or “2006” of the European Working Group of Myelodysplastic Syndromes in Childhood (EWOG-MDS; registered at www.clinicaltrials.gov as #NCT00047268 and #NCT00662090). Approval was obtained from the institutional review board of each institution, and parental informed consent was provided according to the Declaration of Helsinki. Two children were treated at centers not participating in EWOG-MDS studies. One case (D644) was published previously.⁵ The diagnosis of all children was centrally reviewed, and response was evaluated according to international consensus criteria.⁶

The median age of the 12 patients was 4.8 years (range 0.4-9.1) (Table 1). A total of 64 azacitidine cycles were administered (median 5.5 cycles, range 1-11). Seven of 12 treatments consisted of 100 mg/m² azacitidine per intravenous infusion on 5 consecutive days every 28 days. In the other 5 patients, the substance was administered over

5 to 7 days at a single dose of 50 to 100 mg/m² per intravenous or subcutaneous route every 28 to 42 days.

Severe neutropenia ($\leq 500/\mu\text{L}$) was observed in 4 children. Cytopenias led to dose reduction in 2 children, both treated for relapse after second HSCT. Other adverse events were gastrointestinal problems including nausea and vomiting in 2 children, skin rash in 2 children, and fatigue or slight creatinine elevation in 1 patient each. Seven episodes of infection were reported for a total of 64 azacitidine cycles (10.9%).

Of 9 children treated prior to HSCT, 3 normalized blood counts and spleen size (scored as clinical CR) (Table 1). In 2 of these patients, monosomy 7 was present in leukemic cells but disappeared after cycles 5 and 6, respectively. The leukemic karyotype was normal in the other child with clinical CR, precluding the assessment of cytogenetic response. Two of the 3 CR patients featured a somatic *PTPN11* gene mutation. The mutation became undetectable after cycle 6 in 1 child (NS002); material for mutational analysis under azacitidine was unavailable from the other child (D827). The third leukemia carried a somatic *KRAS* mutation, which was no longer detectable after 7 cycles of azacitidine (D644).⁵ One child (CH058) experienced considerable regression of spleen size and became transfusion independent (scored as clinical PR). All 4 children underwent HSCT after 7 to 11 cycles. A fifth child (NS001) responded unusually early, as indicated by reduction of splenomegaly after the first cycle and hematologic improvement after cycle 3 (scored as clinical PR). Azacitidine was then discontinued because of parental choice. Three children progressed rapidly under azacitidine and underwent expedited HSCT. One child (I255) was not evaluable for response because of concomitant treatment. Three children with JMML received azacitidine for leukemia recurrence after the second HSCT. They achieved clinical PR or could be maintained in stable disease for 4 cycles before progressing.

Table 1. Response to azacitidine in children with JMML

Disease status	Patient identifier	Age (y) and gender	Cytogenetics	Mutational group*	Azacitidine cycles	Concomitant treatment	Response to azacitidine†	HSCT (total number)	Status	Follow-up (mo)‡
Prior to first HSCT	A062	1.1, male	Del(5)(q13q33)	<i>NRAS</i>	1 2-3	None None	PR PD	Yes (2)	Alive with leukemia	38
Prior to first HSCT	CH058	2.8, male	Normal	<i>PTPN11</i>	1-3 4-7	None None	SD PR	Yes (2)	Alive with leukemia	16
Prior to first HSCT	D644§	1.4, male	−7	<i>KRAS</i>	1 2-4 5-8	None None None	PD SD CRII	Yes (1)	Alive in remission	62
Prior to first HSCT	D706	5.9, male	−7	<i>PTPN11</i>	1	None	PD	Yes (3)	Alive in remission	66
Prior to first HSCT	D712	6.3, female	Normal	<i>NRAS</i>	1	None	PD	Yes (1)	Dead, TRM	1
Prior to first HSCT	D827	0.4, male	Normal	<i>PTPN11</i>	1-2 3-7	6MP None	Not evaluable CR¶	Yes (1)	Alive in remission	38
Prior to first HSCT	I255	9.1, male	Inv(2)(p23q13), −7	No mutation	1-4	AraC, 6MP	Not evaluable	Yes (1)	Alive in remission	49
Prior to first HSCT	NS001	5.4, male	−7	<i>NRAS</i>	1-3	None	PR	No	Dead, progressive disease	No data
Prior to first HSCT	NS002	0.8, male	−7	<i>PTPN11</i>	1 2-3 4-11	None None None	SD PR CR#	Yes (1)	Dead, TRM	13
Relapse after HSCT**	NL121	4.6, male	Not done	<i>PTPN11</i>	2-4 1, 5-8	None None	SD PD	Yes (3)	Dead, third relapse	6
Relapse after HSCT**	SC108	5.0, male	Normal††	<i>NF1</i>	1-2 3-4 5	None None None	SD PR PD	Yes (2)	Dead, progressive disease	1
Relapse after HSCT**	SC156	5.3, male	Normal††	<i>PTPN11</i>	2-4 1, 5-6	None None	PR PD	Yes (3)	Dead, third relapse	5

AraC, cytarabine; CR, complete response; PD, progressive disease; PR, partial response; SD, stable disease; 6MP, 6-mercaptopurine; TRM, transplantation-related mortality.

*All mutations were confirmed to be somatic. Neurofibromatosis type 1 was diagnosed clinically.

†Cycles of azacitidine with concomitant antineoplastic medication were considered not evaluable.

‡From the end of azacitidine treatment to death or last follow-up.

§This case was published previously by Furlan et al.⁵

¶The patient reached genetic CR with disappearance of monosomy 7 and *KRAS* mutation after 5 and 7 cycles, respectively. Monosomy 7 was tested by fluorescence in situ hybridization, and *KRAS* mutation was assessed by Sanger sequencing.

#Material for mutational studies was not available at the time of clinical CR.

#The patient reached genetic CR with disappearance of monosomy 7 and *PTPN11* mutation after 6 cycles. Monosomy 7 was tested by fluorescence in situ hybridization, and *PTPN11* mutation was assessed by Sanger sequencing.

**Relapse after second HSCT.

††Last cytogenetic analysis: SC108, at start of azacitidine treatment; SC156, prior to second HSCT.

In summary, this retrospective series indicates that low-dose azacitidine is effective and tolerable in JMML and documents 3 cases of JMML where azacitidine induced complete clinical, cytogenetic, and/or molecular genetic remissions before allogeneic HSCT. Importantly, complete remissions without HSCT have not been documented for JMML thus far, regardless of whether conventional cytostatic chemotherapy or newer experimental agents were applied.⁷⁻¹⁰ Prospective clinical investigation of azacitidine, such as the ongoing collaborative Innovative Therapies in Childhood Cancer/EWOG-MDS phase 1/2 trial (Eudra-CT 2010-022235-10), is needed to clarify these questions.

Annamaria Cseh

*Division of Pediatric Hematology-Oncology,
Department of Pediatrics and Adolescent Medicine,
University Medical Center,
Freiburg, Germany*

Charlotte M. Niemeyer

*Division of Pediatric Hematology-Oncology,
Department of Pediatrics and Adolescent Medicine,
University Medical Center,
Freiburg, Germany
German Cancer Consortium (DKTK),
Heidelberg, Germany*

Ayami Yoshimi

*Division of Pediatric Hematology-Oncology,
Department of Pediatrics and Adolescent Medicine,
University Medical Center,
Freiburg, Germany*

Michael Dworzak

*St. Anna Children's Hospital and Children's Cancer Research Institute,
Department of Pediatrics,
Medical University of Vienna,
Vienna, Austria*

Henrik Hasle

*Department of Pediatrics, Aarhus University Hospital,
Skejby, Denmark*

Marry M. van den Heuvel-Eibrink

*Erasmus Medical Centre,
Rotterdam, The Netherlands*

Franco Locatelli

*Department of Pediatric Hematology-Oncology,
Istituto di Ricovero e Cura a Carattere Scientifico (IRCCS) Ospedale Bambino Gesù,
Rome, Italy
University of Pavia,
Pavia, Italy*

Riccardo Masetti

*Pediatric Oncology and Hematology, University of Bologna,
Bologna, Italy*

Markus Schmugge

Department of Hematology and Oncology, University Children's Hospital,
Zurich, Switzerland

Ute Groß-Wieltsch

Olgahospital, Pediatrics 5 (Oncology, Hematology, Immunology,
Rheumatology, Gastroenterology, and General Pediatrics),
Klinikum Stuttgart,
Stuttgart, Germany

Andrea Candás

Department of Hematology and Oncology,
Pediatric Hospital Prof Dr Juan P. Garrahan,
Buenos Aires, Argentina

Andreas E. Kulozik

Department of Pediatric Oncology, Hematology and Immunology,
University of Heidelberg,
Heidelberg, Germany

Lale Olcay

Department of Pediatric Hematology,
Dr Abdurrahman Yurtaslan Ankara Oncology Training and
Research Hospital of the Ministry of Health,
Ankara, Turkey

Meinolf Suttrop

Division of Pediatric Hematology and Oncology, University Children's Hospital,
Dresden, Germany

Ingrid Furlan

University Children's Hospital,
Ulm, Germany

Brigitte Strahm

Division of Pediatric Hematology-Oncology,
Department of Pediatrics and Adolescent Medicine,
University Medical Center,
Freiburg, Germany

Christian Flotho

Division of Pediatric Hematology-Oncology,
Department of Pediatrics and Adolescent Medicine,
University Medical Center,
Freiburg, Germany
German Cancer Consortium (DKTK),
Heidelberg, Germany

Contribution: Annamaria C., C.M.N., and C.F. designed research, analyzed data, and wrote the manuscript; and A.Y., M.D., H.H., M.M.v.d.H.-E., F.L., R.M., M.S., U.G.-W., Andrea C., A.E.K., L.O., M.S., I.F., and B.S. performed research.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

Correspondence: C. Flotho, Division of Pediatric Hematology-Oncology, Department of Pediatrics and Adolescent Medicine, University Medical Center, Mathildenstrasse 1, 79106 Freiburg, Germany; e-mail: christian.flotho@uniklinik-freiburg.de.

References

1. Santini V. Novel therapeutic strategies: hypomethylating agents and beyond. *Hematology Am Soc Hematol Educ Program*. 2012;2012:65-73.
2. Silverman LR, Demakos EP, Peterson BL, et al. Randomized controlled trial of azacitidine in patients with the myelodysplastic syndrome: a study of the cancer and leukemia group B. *J Clin Oncol*. 2002;20(10):2429-2440.
3. Fenaux P, Mufti GJ, Hellstrom-Lindberg E, et al; International Vidaza High-Risk MDS Survival Study Group. Efficacy of azacitidine compared with that of conventional care regimens in the treatment of higher-risk myelodysplastic syndromes: a randomised, open-label, phase III study. *Lancet Oncol*. 2009;10(3):223-232.
4. Gerds AT, Gooley TA, Estey EH, Appelbaum FR, Deeg HJ, Scott BL. Pretransplantation therapy with azacitidine vs induction chemotherapy and posttransplantation outcome in patients with MDS. *Biol Blood Marrow Transplant*. 2012;18(8):1211-1218.
5. Furlan I, Batz C, Flotho C, et al. Intriguing response to azacitidine in a patient with juvenile myelomonocytic leukemia and monosomy 7. *Blood*. 2009;113(12):2867-2868.
6. Niemeyer CM, Loh ML, Cseh A, et al. Criteria for evaluating response and outcome in clinical trials for children with juvenile myelomonocytic leukemia. *Haematologica*. 2015;100(1):17-22.
7. Castleberry RP, Emanuel PD, Zuckerman KS, et al. A pilot study of isotretinoin in the treatment of juvenile chronic myelogenous leukemia. *N Engl J Med*. 1994; 331(25):1680-1684.
8. Niemeyer CM, Arico M, Basso G, et al; European Working Group on Myelodysplastic Syndromes in Childhood (EWOG-MDS). Chronic myelomonocytic leukemia in childhood: a retrospective analysis of 110 cases. *Blood*. 1997;89(10):3534-3543.
9. Castleberry RP, Loh ML, Jayaprakash N, et al. Phase II window study of the farnesyltransferase inhibitor R115777 (Zarnestra®) in untreated juvenile myelomonocytic leukemia (JMML): a children's oncology group study [abstract]. *Blood*. 2005;106(11). Abstract 2587.
10. Bergstraesser E, Hasle H, Rogge T, et al. Non-hematopoietic stem cell transplantation treatment of juvenile myelomonocytic leukemia: a retrospective analysis and definition of response criteria. *Pediatr Blood Cancer*. 2007;49(5):629-633.

Acknowledgments: This work was supported by Deutsche Forschungsgemeinschaft grants FL 345/4-1 (C.M.N., C.F.) and CRC 992-C05 (C.F.).

© 2015 by The American Society of Hematology



blood

2015 125: 2311-2313
doi:10.1182/blood-2015-01-619734

Bridging to transplant with azacitidine in juvenile myelomonocytic leukemia: a retrospective analysis of the EWOG-MDS study group

Annamaria Cseh, Charlotte M. Niemeyer, Ayami Yoshimi, Michael Dworzak, Henrik Hasle, Marry M. van den Heuvel-Eibrink, Franco Locatelli, Riccardo Masetti, Markus Schmugge, Ute Groß-Wieltsch, Andrea Candás, Andreas E. Kulozik, Lale Olcay, Meinolf Suttorp, Ingrid Furlan, Brigitte Strahm and Christian Flotho

Updated information and services can be found at:

<http://www.bloodjournal.org/content/125/14/2311.full.html>

Articles on similar topics can be found in the following Blood collections

[Myeloid Neoplasia](#) (1440 articles)

[Transplantation](#) (2116 articles)

Information about reproducing this article in parts or in its entirety may be found online at:

http://www.bloodjournal.org/site/misc/rights.xhtml#repub_requests

Information about ordering reprints may be found online at:

<http://www.bloodjournal.org/site/misc/rights.xhtml#reprints>

Information about subscriptions and ASH membership may be found online at:

<http://www.bloodjournal.org/site/subscriptions/index.xhtml>